

(+/-1/8"

Tack Seal

4" (+/- 1/8") Micro Vent Holes

3" Customer Seal Area



WITH LAW FOOD 22-0-0

PREVENTS GRABGRASS AND FEEDS IN ONE APPLICATION

LONG-LASTING FEEDING

GROUP 3 HERBICIDE CTIVE INGREDIENT methylpropyl)-6-(trifluoromethyl)-S,S-dimethyl ester..... 0.125 THER INGREDIENTS 99.875 PA Reg. No.62719-483-57131 EPA Est. No. 57131-WI-**KEEP OUT OF REACH OF CHILDREN** CAUTION

Refer to back panel of bag for Directions for Use

Tack Seal 1.5"

(+/-1/8'')

From Top Of Bag

APPLICATION EARLY SPRING

SEASON-LONG CRABGRASS CONTROL

RICH GREEN LAWN

GOVERS 5,000 Sq. Ft. NET WEIGHT 16 LBS. (7.26 KG)



TOTAL NITROGEN (N) 22.0% Urea Nitrogen* SOLUBLE POTASH (K2O). IRON (Fe)..

mary Plant Foods Derived from: Urea, pated Sulfur Coated Urea and Muriate of Potasi on derived from Iron Oxide and Ferrous Sulfate 4.4% slowly available urea nitrogen from po



BROADCAST S	PREADER	DROP SPREADER		
Brand/Model of Spreader	Calibration Settings at 3.2 lb./1,000 sq.ft.	Brand/Model of Spreader	Calibration Settings at 3.2 lb./1,000 sq.ft.	
AgriFab	4	AgriFab	4	
Craftsman	4	Lawncrafters	4	
Earthway EV-N-Spread	12	Craftsman	4	
Precision	4	Earthway EV-N-Spread	9.5	
Red Devil	4	Precision	1.5	
Scotts Basic	4.5	Red Devil	1.5	
Scotts Standard 4.5		Scotts AccuGreen 1000	6	
Scotts Deluxe EdgeGuard 4.5		Scotts AccuGreen 2000 6		
Scotts Easy Green 26		Scotts AccuGreen 3000	6	
Scotts Speedy Green	5.75	Scotts Even Green	7.5	

MAXLAWN Maximize Your Lawn Crabgrass Preventer with Lawn Food For control of crabgrass and control or suppression of other listed annual grass and broadleaf weeds in established lawns and ornamental turf. Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes. Not for use on plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes. For use on plants intended for aesthetic purposes or climatic modification and being grown on golf courses or lawns. In the state of New York, this product may be applied only by commercial applicators at aminimum of 200 lb (0.25 lb active ingredient) per acre (9.2 lb per 1000 sq ft) per year. Use of this dithiopyr is a Group 3 herbicide based on the mode of action

product in Nassau and Suffolk counties of New York is prohibited. It is a violation of law in Maryland and New Jersey to apply this product to turf at a rate that results in greater than 0.7 lb per 1,000 sq.ft. of readily available nitrogen or more than 0.9 lb per 1,000 sq.ft. of total nitrogen with a least 20% of which shall consist of slow release nitrogen per application. Additional restrictions apply to total nitrogen applied to turf per season. Other state and local restrictions ma apply. [Maryland or New Jersey Applications]

Notice: Read the entire label. Use only according to label directions. Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies at end of Directions for Use. If terms are unacceptable, return at once unopened. In case of emergency endangering health or the environment involving this product, call 1-800-992-5994. Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

Crab-Buster is a Registered Trademark of The Dow Chemical Company ("Dow") or an affiliated company of Dow Dow AgroSciences LLC • Indianapolis, IN 46268 U.S.A.

If in eyes: Hold eye open and rinse slowly and gently with water | This product should be applied only to lawns and ornamental for 15-20 minutes. Remove contact lenses, if present, after the turfgrasses that are well established. For newly established lawns, first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

initial application of this product should be made only after turfgrasses have developed a vigorous root system, uniform stand,

center or doctor for further treatment advice.

Hot Line Number: Have the product container or label with you

• Use of this product on lawns and ornamental turfgrasses that are when calling a poison control center or doctor, or going for not well established, or weakened by stress due to mechanical treatment. You may also contact 1-800-992-5994 day or night, operations, pests, excessive temperatures, or lack of moisture may for emergency treatment information.

PRECALITIONARY STATEMENTS HAZARDS TO HUMANS & DOMESTIC ANIMALS CAUTION: Causes eye irritation • Harmful if inhaled Avoid contact with eyes or clothing. Avoid breathing dust. Wash these cultural practices.

cause allergic reaction in some individuals. Environmental Hazards

This product is toxic to fish and highly toxic to other aquatic cultural practices for seedbed preparation, seeding, irrigation and organisms including oysters and shrimp. To protect the fertilization. environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this Use Restrictions that lands on a driveway, sidewalk, or street, back onto the settled. treated area of the lawn or garden will help to prevent runoff to water bodies or drainage systems.

Directions for Use inconsistent with its labeling.

Storage and Disposal
Do not contaminate water, foodstuffs, feed or seed by storage or
Dimension® 0.1% FG turf and ornamental herbicide as single or
Dimension® 0.1% FG turf and ornamental herbicide as single or
Dimension® 0.1% FG turf and ornamental herbicide as single or
Dimension® 0.1% FG turf and ornamental herbicide as single or
Dimension® 0.1% FG turf and ornamental herbicide as single or
Dimension® 0.1% FG turf and ornamental herbicide as single or
Dimension® 0.1% FG turf and ornamental herbicide as single or
Dimension® 0.1% FG turf and ornamental herbicide as single or
Dimension® 0.1% FG turf and ornamental herbicide as single or
Dimension® 0.1% FG turf and ornamental herbicide as single or
Dimension® 0.1% FG turf and ornamental herbicide as single or
Dimension® 0.1% FG turf and ornamental herbicide as single or
Dimension® 0.1% FG turf and ornamental herbicide as single or
Dimension® 0.1% FG turf and ornamental herbicide as single or
Dimension® 0.1% FG turf and ornamental herbicide as single or
Dimension® 0.1% FG turf and ornamental herbicide as single or
Dimension® 0.1% FG turf and ornamental herbicide as single or
Dimension® 0.1% FG turf and ornamental herbicide as single or
Dimension® 0.1% FG turf and ornamental herbicide as single or
Dimension® 0.1% FG turf and ornamental herbicide as single or
Dimension® 0.1% FG turf and ornamental herbicide as single or
Dimension® 0.1% FG turf and ornamental herbicide as single or
Dimension® 0.1% FG turf and ornamental herbicide as single or
Dimension® 0.1% FG turf and ornamental herbicide as single or
Dimension® 0.1% FG turf and ornamental herbicide as single ornamental Pesticide Storage: Store this product only in its original ingredient) per acre. Applicators may choose to make a single application or sequential applications at 200 to 400 pounds per acre Container Handling: Nonrefillable container. Do not reuse or at 5 to 10 week intervals based on one or more of the factors listed refill this container. **If empty:** Place in trash or offer for recycling if available. If partially filled: Call your local solid waste agency or 1-800-CLEANUP for disposal instructions. Never place unused

Use Directions for Turf This turf fertilizer product, containing Crab-Buster® herbicide, can be applied up to 4 weeks later than other crabgrass preventers and provides up to 4 months of crabgrass control. MAXLAWN Maximize Your Lawn Crabgrass Preventer with Lawn Food provides preemergence control of crabgrass and control or suppression of other listed weeds when applied prior postemergence control of seedling crabgrass after germination and emergence. Postemergence control is limited to crabgrass

Tips for Improved Weed Control

• For best results, apply this product within a few days after mowing only, but is effective up to 4 weeks after crabgrass has and delay mowing again for a few days after the application. crabgrass germination. Apply this product only to turfgrass species listed on this label. irrigation. Erratic weed control may result if or irrigation within 30 days after application. The following turfgrass species are tolerant to this product when Weeds Controlled or Suppressed

Cool-Season Grasses Warm-Season Grasses

Bentgrass, creeping Bluegrass, Kentlucky Fescue, fine Personal Responsibility of the Control of Crabgrass and Control of Crabgras

Use Precautions for Turf

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice.

Thinhaled: Move person to fresh air. If person is not breathing, and received at least two mowings following seeding, sodding, or sprigging. Exception: In newly established lawns of Kentucky bluegrass, perennial ryegrass and/or tall fescue, this product may be applied two weeks after the first sign of germination.

 For best results, cultural practices that disturb the soil, such as core-, spike-, or hydro-aerification, and verticutting, should be done before application of this product. Do not apply this product until turfgrass has recovered from the mechanical injury caused by thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse. Prolonged or frequently repeated skin contact while handling this material may cause allernic reaction in some individuals. application program totaling 9.2 lb/1000 sq ft (0.5 lb active ingredient per acre) or more may inhibit the establishment of desirable turfgrasses. When reseeding, overseeding follow proper

product in calm weather when rain is not predicted for the next

• Do not use clippings from treated turf for mulching around 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Sweeping any product

vegetables or fruit trees.

• Do not allow people or pets to enter treated area until dust has

Application Directions for Turf Apply Dimension® 0.125% FG turf and ornamental herbicide with equipment that provides a uniform particle distribution. Calibrate It is a violation of federal law to use this product in a manner application equipment prior to usage. Avoid streaking, skips, or excess overlaps during application.

> Preemergence Application Rates, Frequency and Timing sequential application at 200 to 400 pounds (0.25 to 0.5 lb active below. Avoid streaking, skips, or excess overlaps during application. I enoth of residual weed control desired Height of turf (lower cut turf may require higher use rates)

• History and success of weed control at the application site (use higher listed application rates if herbicide treatment history is unknown or weed control was poor with previous applications) Exposure to high temperatures and heavy rainfall or irrigation (this will shorten the residual preemergence performance) On turf sites adjacent to hard surfaces including driveways. sidewalks and parking lots where residual activity may be annua, goosegrass and sandbur) will require higher use rates

and delay mowing again for a few days after the application.

This generally corresponds to the time when crabgrass seedlings first become visible in the established turfgrass. Do not apply this product later than 4 weeks after than 4 weeks after than 4 weeks after the application and delay mowing again for a few days after the application.

The turf herbicide and fertilizer components in this product are not effective until activated by rainfall or sprinkled turfgrass. Do not apply this product later than 4 weeks after the application.

The turf herbicide and fertilizer components in this product are not effective until activated by rainfall or sprinkled turfgrass. Do not apply this product later than 4 weeks after the application.

The turf herbicide and fertilizer components in this product are not effective until activated by rainfall or sprinkled turfgrass. Do not apply this product later than 4 weeks after the application.

The turf herbicide and fertilizer components in this product are not effective until activated by rainfall or irrigation. Performance is interest the application and delay mowing again for a few days after the application.

The turf herbicide and fertilizer components in this product are not effective until activated by rainfall or irrigation. Performance is interest the application and delay mowing again for a few days after the application.

The turf herbicide and fertilizer components in this product are not effective until activated by rainfall or irrigation. Performance is not effective until activated by rainfall or irrigation. Performance is not effective until activated by rainfall or irrigation. Performance is not effective until activated by rainfall or irrigation. Performance is not effective until activated by rainfall or irrigation. Performance is not effective until activated by rainfall or irrigation. Performance is not effective until activated by rainfall or irrigation. Performance is not effective until activated by rainfall or irrigation. Performance is not effective

MAXLAWN Maximize Your Lawn Crabgrass Preventer with Lawn is being applied. Refer to the map below to determine the Food will provide control of crabgrass and control or suppression of

GUARANTEED ANALYSI

	Grasses	Broadleaf Weeds		Use	
	barley barnyardgrass bluegrass, annual brome crabgrass, large crabgrass, smooth crabgrass, Southern	bittercress † carpetweed chickweed † geranium, Carolina † henbit knotweed, prostrate lespedeza, common †	speedwell, corn † spurge, garden spurge, prostrate spurge, spotted woodsorrel, creeping woodsorrel, yellow		
	crowfootgrass † dallisgrass	marestail medic, black		Use	
	goosegrass foxtail, green	mustard oxalis, buttercup		(Po	
	foxtail, yellow kikuyugrass † oats, wild	pineappleweed pigweed, redroot parsley-piert †		ss Control	
	niegrass	nurslane common		ll ss	

classification system of the Weed Science Society of America. herbicides is a best practice. A diversified weed management Maximum Application Rates program may include the use of multiple herbicides with different • Do not apply more than 9.2 lb of this product per 1,000 sq ft per modes of action and overlapping weed spectrum with or without application. tillage operations and/or other cultural practices. Research has • Do not apply more than 27.6 lb of this product/1000 sq ft per demonstrated that using the labeled rate and directions for use is year (equivalent to 0.5 and 1.5 lb of active ingredient per acre,

important to delay the selection for resistance.

To aid in the prevention of developing weeds resistant to this. This product may be applied with drop or rotary-type spreaders product, users should:

will be appropriate for the weed species and weed sizes Start with a clean field, using either a burndown herbicide application or tillage. • If using post-emergence herbicides or tank mixes, control weeds early when they are relatively small. Apply full rates of Dimension 0.125% FG for the most difficult this product uniformly over the treatment area. More uniformity

weed escapes. in control of weed species. Control weed escapes before they reproduce by seed or proliferate vegetatively. Report any incidence of non-performance of this product against a particular weed to your local company representative,

Terms and Conditions of Use

local retailer, or county extension agent. Contact your local company representative, crop advisor, or of this product specified for your local conditions. Tank mix products so that there are multiple effective modes of action • If resistance is suspected, treat weed escapes with an herbicide having a mode of action other than Group 3 and/or

use nonchemical methods to remove escapes, as practical, with the goal of preventing further seed production. Suspected herbicide-resistant weeds may be identified by Failure to control a weed species normally controlled by

the herbicide at the dose applied, especially if control is achieved on adjacent weeds; A spreading patch of non-controlled plants of a particular weed species; and Surviving plants mixed with controlled individuals of the

herbicide resistance management practices as is practical: Use a broad spectrum herbicide with other mode of action as a foundation in a weed control program, if appropriate. Utilize sequential applications of herbicides with alternative modes of action. Rotate the use of this product with non-Group 3 herbicides.

 Avoid making more than two sequential applications of Dimension 0.125% FG and any other Group 3 herbicides within a single growing season unless mixed with an herbicide with a different mode of action with an overlapping spectrum for the difficult-to-control weeds. Incorporate non-chemical weed control practices, such as Use good agronomic principles that enhance crop

development and crop competitiveness. • Thoroughly clean plant residues from equipment before I eaving fields suspected to contain resistant weeds. Manage weeds in and around fields to reduce weed seed

irrigation. Erratic weed control may result if not activated by rainfall control will vary depending on use rate, weather conditions, condition of the turfgrass, intensity of weed pressure, and the location or "Region" within the United States where this product **region of the U.S. that includes your location.** Use the table

"Regions" within the United States

2.4 lb/1000 sq ft 0.128 lb ai/acre) 3.2 lb/1000 sq ft 4.0 - 4.7 lb/1000 sq ft † .174 lb ai/acre) .0 - 9.2 lb/1000 sg ft + 4.0 - 9.2 lb/1000 sg ft +

1,000 square feet equals an area 100 feet x 10 feet or square area of out 32 feet on a side.

Jse the higher rate in rate range in areas of high weed pressure or for reased length of control.

The continued effectiveness of this product depends on the successful implementation of a weed resistance management year of this product (equivalent to 0.25 lb of active ingredient per program

Application Instructions and Equipment designed to apply granular herbicides. For best results, apply • Scout fields before application to ensure herbicides and rates this product evenly and uniformly avoiding streaking, skips or overlaps. Avoid the use of spreaders that tend to apply granules according to the manufacturer's directions. Initial spreader settings may require adjustment to deliver the application rate setting may be marked or recorded for future reference. Apply to control weed in the field at the specified time to minimize of application can usually be achieved by applying one-half of the required amount of product over the treatment area and then • Scout fields after application to detect weed escapes or shifts applying the remaining one-half in a different direction (e.g., at a right angle to the previous direction). Avoid streaking, skips, or overlaps during application. Check equipment frequently to verify calibration and proper

If terms of the following Warranty Disclaimer, Inherent Risks of Use, and Contact your local company representative, crop advisor, or extension agent to find out if suspected resistant weeds to this MOA have been found in your region. If resistant biotypes
 Limitation of Remedies are not acceptable, return unopened package at once to the seller for a full refund of purchase price paid. To the extent permitted by law, otherwise, use by the buyer or any other user constitutes of target weeds have been reported, use the application rates acceptance of the terms under Warranty Disclaimer, Inherent Risks of

> Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. To the extent permitted by law, SELLER MAKES NC OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER Inherent Risks of Use

is impossible to eliminate all risks associated with use of this product Plant injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperature, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of Seller. To the extent permitted by law, all such risks

damages resulting from this product (including claims based on contra regligence, strict liability, or other legal theories), shall be limited to, at Seller's election, one of the following: . Refund of purchase price paid by buyer or user for

product bought, or 2. Replacement of amount of product used. To the extent permitted by law, seller shall not be liable for losses or damages resulting from handling or use of this product unless Seller is promptly notified of such loss or damage in writing. To the extent permitted by law, in no case shall Seller be liable for consequential or

The terms of the Warranty Disclaimer, Inherent Risks of Use, and this mechanical cultivation, crop rotation, cover crops and weedfree crop seeds, as part of an integrated weed control program. statements or agreements. No employee or sales agent of the Seller or the seller is authorized to vary or exceed the terms of the Warranty Disclaimer or this Limitation of Remedies in any manner.

> FPO UPC 0 24378 61102 6

Doc.# RP62719-080-18-16

DISTRIBUTED AND GUARANTEED BY EAU CLAIRE COOP OIL CO., P.O. Box 837, Eau Claire, WI 54702

Prepared By Direct Plastics Sept 23 2011 Max Lawn 16lb Lawn Food 26-0-4 (720-2020B

Max Lawn 16lb Winterizer 22-0-10 (720-2030B) Max Lawn 16lb CrabGrass 22-0-4 (720-2000B) Max Lawn 16lb Starter 12-24-6 (720-2040B)